

# Amphilophium buccinatorium

## COMMON NAME

Cherere

## FAMILY

Bignoniaceae

## AUTHORITY

Amphilophium buccinatorium (DC.) L.G.Lohmann

## FLORA CATEGORY

Vascular – Exotic

## STRUCTURAL CLASS

Lianes & Related Trailing Plants - Dicotyledons

## CONSERVATION STATUS

Not applicable

## SIMILAR TAXA

Previously known as Distictis buccinatoria and Phaedoranthus buccinatoria

## FLOWER COLOURS

Orange, Red/Pink

## YEAR NATURALISED

1988

## ORIGIN

Probably Mexico

## Reason For Introduction

Horticultural

## Life Cycle Comments

Long-lived woody perennial vine

## Reproduction

Vegetative spread through layering

## Seed

Not formed in New Zealand

## Dispersal

Vegetative spread, usually from carelessly discarded garden waste

## REFERENCES AND FURTHER READING

Lohmann, L.G.; Taylor, C.M. 2014: A new generic classification of tribe Bignoniaceae. *Annals of the Missouri Botanical Garden* 99(3): 348–489.

## NZPCN FACT SHEET CITATION


Please cite as: Ogle, C. (Year at time of access): *Amphilophium buccinatorium* Fact Sheet (content continuously updated). New Zealand Plant Conservation Network.

<https://www.nzpcn.org.nz/flora/species/amphilophium-buccinatorium/>

(Date website was queried)

## MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/amphilophium-buccinatorium/>



In cultivation, Auckland. Nov 2006.

Photographer: Peter J. de Lange, Licence: CC BY-NC.



In cultivation, Auckland. Nov 2006.

Photographer: Peter J. de Lange, Licence: CC BY-NC.