

Anisotome aromatica

COMMON NAME

aromatic aniseed, kopoti, common aniseed

FAMILY

Apiaceae

AUTHORITY

Anisotome aromatica Hook.f.

FLORA CATEGORY

Vascular – Native

ENDEMIC TAXON

Yes

ENDEMIC GENUS

No

ENDEMIC FAMILY

No

STRUCTURAL CLASS

Herbs - Dicotyledons other than Composites

NVS CODE

ANIARO

CHROMOSOME NUMBER

2n = 22

CURRENT CONSERVATION STATUS

2017 | Not Threatened

PREVIOUS CONSERVATION STATUSES

2012 | Not Threatened

2009 | Not Threatened

2004 | Not Threatened

WETLAND PLANT INDICATOR STATUS RATING

FACU: Facultative Upland

Occasionally is a hydrophyte but usually occurs in uplands (non-wetlands).

FLOWER COLOURS

White

LIFE CYCLE

Winged mericarps are dispersed by wind (Thorsen et al., 2009).

ETYMOLOGY

anisotome: Unequal sided

aromatica: Aromatic

REFERENCES AND FURTHER READING

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. Perspectives in Plant Ecology, Evolution and Systematics 2009 Vol. 11 No. 4 pp. 285-309

MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/anisotome-aromatica/>



Mangatepopo, November. Photographer: John Smith-Dodsworth, Licence: CC BY-NC.



Anisotome aromatica. Photographer: Jane Gosden, Date taken: 28/11/2009, Licence: CC BY-NC-SA.