

# Anisotome capillifolia

## FAMILY

Apiaceae

## AUTHORITY

Anisotome capillifolia (Cheeseman) Cockayne

## FLORA CATEGORY

Vascular – Native

## ENDEMIC TAXON

Yes

## ENDEMIC GENUS

No

## ENDEMIC FAMILY

No

## STRUCTURAL CLASS

Herbs - Dicotyledons other than Composites

## NVS CODE

ANICAP

## CHROMOSOME NUMBER

2n = 22

## CURRENT CONSERVATION STATUS

2017 | At Risk – Declining | Qualifiers: DP, PD, RF

## PREVIOUS CONSERVATION STATUSES

2012 | Not Threatened

2009 | Not Threatened

2004 | Not Threatened

## FLOWER COLOURS

White

## LIFE CYCLE

Winged mericarps are dispersed by wind (Thorsen et al., 2009).

## ETYMOLOGY

**anisotome:** Unequal sided

**capillifolia:** Hairy leaved

## REFERENCES AND FURTHER READING

Thorsen MJ, Dickinson KJM, Seddon PJ. 2009. Seed dispersal systems in the New Zealand flora. *Perspectives in Plant Ecology, Evolution and Systematics* 11: 285–309. <https://doi.org/10.1016/j.ppees.2009.06.001>.

## MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/anisotome-capillifolia/>



Mt Burns, January. Photographer: John Smith-Dodsworth, Licence: CC BY-NC.



Gertrude Valley. Photographer: John Barkla, Licence: CC BY.