

Azorella cockaynei

SYNONYMS

Schizeilema cockaynei (Diels) Cheeseman

FAMILY

Apiaceae

AUTHORITY

Azorella cockaynei Diels

FLORA CATEGORY

Vascular – Native

ENDEMIC TAXON

Yes

ENDEMIC GENUS

No

ENDEMIC FAMILY

No

STRUCTURAL CLASS

Herbs - Dicotyledons other than Composites

NVS CODE

SCHCOC

CHROMOSOME NUMBER

2n = 48

CURRENT CONSERVATION STATUS

2017 | Not Threatened

PREVIOUS CONSERVATION STATUSES

2012 | Not Threatened

2009 | Not Threatened

2004 | Not Threatened

DISTRIBUTION

Endemic. South and Stewart Islands, local from Otago south throughout Southland and on Stewart island

WETLAND PLANT INDICATOR STATUS RATING

FACW: Facultative Wetland

Usually is a hydrophyte but occasionally found in uplands (non-wetlands).

FLOWERING

November - December

FRUITING

December - March

PROPAGATION TECHNIQUE

Easily grown in a damp soil. Best within a partially submerged pot.

ETYMOLOGY

cockaynei: Named after Leonard Cockayne FRS (7 April 1855 - 8 July 1934) who is regarded as New Zealand's greatest botanist and a founder of modern science in New Zealand



Azorella cockaynei in alpine streamside herbfield, Old Man Range. Photographer: John Barkla, Date taken: 21/03/2020, Licence: CC BY.



Azorella cockaynei in coastal turf with Samolus repens, Riverton, Southland. Photographer: John Barkla, Date taken: 25/12/2017, Licence: CC BY.

WHERE TO BUY

Not commercially available.

MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/azorella-cockaynei/>