

# Azorella haastii var. haastii

## SYNONYMS

Pozoa haastii Hook.f., Pozoa elegans Colenso, Azorella haastii (Hook.f.) Kirk, Schizeilema haastii subsp. hookerianum Domin, Schizeilema haastii (Hook.f.) Domin var. haastii

## FAMILY

Apiaceae

## AUTHORITY

Azorella haastii (Hook.f.) Drude var. haastii

## FLORA CATEGORY

Vascular – Native

## ENDEMIC TAXON

Yes

## ENDEMIC GENUS

No

## ENDEMIC FAMILY

No

## STRUCTURAL CLASS

Herbs - Dicotyledons other than Composites

## NVS CODE

SCHHVH

## CHROMOSOME NUMBER

2n = 32

## CURRENT CONSERVATION STATUS

2017 | Not Threatened

## PREVIOUS CONSERVATION STATUSES

2012 | Not Threatened

2009 | Not Threatened

2004 | Not Threatened

## DISTRIBUTION

Endemic. North and South Islands from the Ruahine Range south.

## FLOWERING

December - February

## FRUITING

January - April

## ETYMOLOGY

**haastii**: Honours the New Zealand geologist and botanist Sir Julius von Haast (1822-87)

## REFERENCES AND FURTHER READING

Plunkett, G.M.; Nicolas, A.N. 2016: Assessing *Azorella* (Apiaceae) and its allies: Phylogenetics and a new classification. *Brittonia* 68:1-31.

## MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/azorella-haastii-var-haastii/>



Homer Basin, January. Photographer: John Smith-Dodsworth, Licence: CC BY-NC.