# **Buellia disciformis**

#### **SYNONYMS**

Lecidea parasema var. disciformis Fr., Hafellia disciformis (Fr.) Marbach & H.Mayrhofer

#### **FAMILY**

Caliciaceae

#### **AUTHORITY**

Buellia disciformis (Fr.) Mudd

### **FLORA CATEGORY**

Lichen - Native

## **ENDEMIC TAXON**

Nο

## **ENDEMIC GENUS**

Nο

#### **ENDEMIC FAMILY**

No

## STRUCTURAL CLASS

Lichens - Crustose

## **CURRENT CONSERVATION STATUS**

2018 | Not Threatened | Qualifiers: SO

## **BRIEF DESCRIPTION**

Characterised by the corticolous habit; the thin, grey-white thallus; sessile, black apothecia; and ascospores with slight apical thickenings (*Callispora*-type),  $14-30 \times 6-13 \mu m$ .

## **DISTRIBUTION**

**South Island:** Canterbury (Arthur's Pass, Hanmer, Lake Lyndon), Southland (South Mavora Lake).

Known also from Great Britain, Europe, Scandinavia, Canary Islands, Socotra, North America and Australia.

#### **HABITAT**

On bark of native and introduced trees, \*Alnus glutinosa, Dracophyllum acerosum, Fuscospora cliffortioides. Often associating on mountain beech bark with the lichens Haematomma alpinum, Miltidea ceroplasta, Ochrolechia etc.

### **DETAILED DESCRIPTION**

**Thallus** thin, areolate, white to grey, irregular. **Medulla** ±amyloid. **Apothecia** sessile, to 1.5 mm diam., disc black, epruinose, margins usually persistent. **Hymenium** with oil droplets; epithecium olive-brown to brown. **Hypothecium** dark-brown. **Ascospores** *Callispora*-type, 1-septate, rarely 3-septate, apices somewhat pointed,  $14-30 \times 6-13 \mu m$ . **Pycnidia** common, conidia subcylindrical,  $4-5 \times 1 \mu m$ .

**Chemistry:** Thallus K+ yellow; containing atranorin.

## **SUBSTRATE**

Corticolous





Image: taken by A.J. Marshall based on herbarium material collected by P.J. de Lange from the canopy branchlets of Metrosideros excelsa × M. robusta. Te Paki, Unuwhao Forest, below Tarure Hill. Photographer: Peter J. de Lange, Date taken: 20/06/2023, Licence: CC BY.



Image: taken by A.J. Marshall based on herbarium material collected by P.J. de Lange from the canopy branchlets of Metrosideros excelsa × M. robusta. Te Paki, Unuwhao Forest, below Tarure Hill. Photographer: Peter J. de Lange, Date taken: 20/06/2023, Licence: CC BY.

#### **ATTRIBUTION**

Fact sheet prepared by Marley Ford (12 February 2021). Brief description, Distribution, Habitat, and Features sections copied from Galloway (1985) & Galloway (2007).

# REFERENCES AND FURTHER READING

Galloway D.J. 2007: Flora of New Zealand: Lichens, including lichen-forming and lichenicolous fungi. 2nd edition. Lincoln, Manaaki Whenua Press. 2261 pp.

## **MORE INFORMATION**

https://www.nzpcn.org.nz/flora/species/buellia-disciformis/