

# Centrolepis glabra

## FAMILY

Restionaceae

## AUTHORITY

*Centrolepis glabra* (F.Muell. ex Sond.) Hieron

## FLORA CATEGORY

Vascular – Native

## ENDEMIC TAXON

No

## ENDEMIC GENUS

No

## ENDEMIC FAMILY

No

## STRUCTURAL CLASS

Rushes & Allied Plants

## CURRENT CONSERVATION STATUS

2017 | At Risk – Naturally Uncommon | Qualifiers: DP, SO, Sp

## DISTRIBUTION

Indigenous. New Zealand: South Island from South Canterbury to Fiordland.

## ETYMOLOGY

**centrolepis:** Pointed scale

**glabra:** Hairless

## TAXONOMIC NOTES

*Centrolepis* was formerly in the family Centrolepidaceae, which was shown by Briggs et al. (2014) to be embedded within Restionaceae. Based on these findings, the Angiosperm Phylogeny Group adopted an enlarged concept of Restionaceae that included Centrolepidaceae (APG IV 2016), which is followed by NZPCN.

## REFERENCES AND FURTHER READING

Angiosperm Phylogeny Group. 2016. An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG IV. *Botanical Journal of the Linnean Society* 181: 1–20. <https://doi.org/10.1111/boj.12385>.

Briggs BG, Marchant AD, Perkins AJ. 2014. Phylogeny of the restiid clade (Poales) and implications for the classification of Anarthriaceae, Centrolepidaceae and Australian Restionaceae. *Taxon* 63: 24–46. <https://doi.org/10.12705/631.1>.

## MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/centrolepis-glabra/>



Hakatere Conservation Area. Photographer: Jane Gosden, Date taken: 21/02/2024, Licence: CC BY-NC-SA.



Hakatere Conservation Area. Photographer: Jane Gosden, Date taken: 21/02/2024, Licence: CC BY-NC-SA.