

Eremophila debilis

FAMILY

Scrophulariaceae

AUTHORITY

Eremophila debilis (Andrews) Chinnock

FLORA CATEGORY

Vascular – Exotic

STRUCTURAL CLASS

Trees & Shrubs - Dicotyledons

CONSERVATION STATUS

Not applicable

FLOWER COLOURS

Red/Pink, White

LIFE CYCLE

Fleshy berries are dispersed by frugivory (Thorsen et al., 2009).

ORIGIN

Australia. Long regarded as indigenous to New Zealand, where it has been known as *Myoporum debile*.

REASON FOR INTRODUCTION

Probably horticultural

ETYMOLOGY


debilis: Weak

REFERENCES AND FURTHER READING

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. *Perspectives in Plant Ecology, Evolution and Systematics* 11: 285-309

MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/eremophila-debilis/>



Naturalised at Oratia. Photographer: Jeremy R. Rolfe, Date taken: 19/07/2007, Licence: CC BY.



Naturalised at Oratia. Photographer: Jeremy R. Rolfe, Date taken: 19/07/2007, Licence: CC BY.