

Euchiton lateralis

SYNONYMS

Gnaphalium laterale C.J.Webb

FAMILY

Asteraceae

AUTHORITY

Euchiton lateralis (C.J.Webb) Breitw. et J.M.Ward

FLORA CATEGORY

Vascular – Native

ENDEMIC TAXON

Indeterminate

ENDEMIC GENUS

No

ENDEMIC FAMILY

No

STRUCTURAL CLASS

Herbs - Dicotyledonous composites

NVS CODE

EUCLAT

CHROMOSOME NUMBER

2n = 28

CURRENT CONSERVATION STATUS

2017 | Not Threatened

PREVIOUS CONSERVATION STATUSES

2012 | Not Threatened

2009 | Not Threatened

2004 | Not Threatened

WETLAND PLANT INDICATOR STATUS RATING

FACW: Facultative Wetland

Usually is a hydrophyte but occasionally found in uplands (non-wetlands).

LIFE CYCLE

Pappate cypselae are dispersed by wind and water (Thorsen et al., 2009).

ETYMOLOGY

euchiton: From the Greek eu (good) and chiton (tunic or covering)

REFERENCES AND FURTHER READING

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. Perspectives in Plant Ecology, Evolution and Systematics 11: 285-309

MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/euchiton-lateralis/>

