

Forstera sedifolia

SYNONYMS

Forstera glabra Gaertn.; *Forstera sedifolia* var. *oculata* Cheeseman

FAMILY

Stylidiaceae

AUTHORITY

Forstera sedifolia G.Forst.

FLORA CATEGORY

Vascular – Native

ENDEMIC TAXON

Yes

ENDEMIC GENUS

No

ENDEMIC FAMILY

No

STRUCTURAL CLASS

Herbs - Dicotyledons other than Composites

CURRENT CONSERVATION STATUS

2017 | Not Threatened

PREVIOUS CONSERVATION STATUSES

2012 | Not Threatened

2009 | Not Threatened

2004 | Not Threatened

WETLAND PLANT INDICATOR STATUS RATING

FACU: Facultative Upland

Occasionally is a hydrophyte but usually occurs in uplands (non-wetlands).

FLOWER COLOURS

White

LIFE CYCLE

Winged seeds are dispersed by wind (Thorsen et al., 2009).

ETYMOLOGY

forstera: Named after Johann Reinhold Forster and his son Johann Georg Adam Forster who were 18th century naturalists on James Cook's second voyage

sedifolia: Sedum-leaved

REFERENCES AND FURTHER READING


Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. *Perspectives in Plant Ecology, Evolution and Systematics* 11: 285-309

MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/forstera-sedifolia/>



Mt Arthur February. Photographer: John Smith-Dodsworth, Licence: CC BY-NC.



Kepler Mountains, Fiordland. Photographer: Jesse Bythell, Licence: CC BY-NC.