Lecanora argentata

SYNONYMS

Parmelia subfusca f. ("n") argentata Ach.

FAMILY

Lecanoraceae

AUTHORITY

Lecanora argentata (Ach.) Degel.

FLORA CATEGORY

Lichen - Native

ENDEMIC TAXON

No

ENDEMIC GENUS

Nο

ENDEMIC FAMILY

Νo

STRUCTURAL CLASS

Lichens - Crustose

CURRENT CONSERVATION STATUS

2018 | At Risk - Naturally Uncommon | Qualifiers: DP, SO

BRIEF DESCRIPTION

Recognised by the corticolous habit; the clear, red-brown epithecium; large crystals in the amphithecium; and products of the gangaleoidin chemosyndrome.

DISTRIBUTION

North Island: Northland (Little Barrier Island, Waiwera) to Auckland (Mt Albert Research Station).

Known from Great Britain, Europe, Scandinavia and North America.

HABITAT

In coastal forest (*Metrosideros*) and on a planted specimen of *Fuscospora*. It occurs in high-light habitats with species of *Buellia*, *Flavoparmelia*, *Pertusaria* and *Parmotrema*.

DETAILED DESCRIPTION

Thallus continuous, yellowish to greenish white, or greyish smooth at margins, uneven to subverrucose centrally, without pruina or soredia, delimited by a \pm prominent grey-white or whitish prothallus. **Apothecia** sessile, 0.4–1 mm diam., scattered or crowded centrally; thalline exciple persistent, concolorus with thallus, smooth to \pm verrucose or crenulate; discs reddish or dark red-brown to bright red-brown, epruinose. **Amphithecium** with large crystals (pulicaris-type not dissolving in K); epithecium reddish brown not dissolving in K, without crystals (glabrata-type), 10–15 μ m thick. **Paraphyses** c. 2 μ m thick, simple, dense, apices capitate to 4 μ m thick. **Hymenium** colourless, 55–70 μ m tall. **Hypothecium** colourless. **Asci** clavate 40–55 × 10–14 μ m. **Ascospores** ellipsoidal, 10.5–15.5 × 5.5–8.5 μ m, the wall 0.5 μ m thick.

Chemistry: Thallus and apothecial margin K+ yellow, C-, KC-, Pd+ yellow; containing atranorin and gangaleoidin as major compounds with chloroatranorin, 4-dechlorogangaleoidin and norgangaleoidin as minor compounds

SUBSTRATE

Corticolous





Local on London plane branch, Mt Albert Auckland. Photographer: Marley Ford, Date taken: 10/09/2021, Licence: CC BY.



Local on London plane branch, Mt Albert Auckland. Photographer: Marley Ford, Date taken: 10/09/2021, Licence: CC BY.

ATTRIBUTION

Fact sheet prepared by Marley Ford (3 August 2022). Brief description, Distribution, Habitat, and Features sections copied from Galloway (2007).

REFERENCES AND FURTHER READING

Galloway D.J. 2007: Flora of New Zealand: Lichens, including lichen-forming and lichenicolous fungi. 2nd edition. Lincoln, Manaaki Whenua Press. 2261 pp.

MORE INFORMATION

https://www.nzpcn.org.nz/flora/species/lecanora-argentata/