# **Placopsis lateritioides**

# **FAMILY**

Trapeliaceae

#### **AUTHORITY**

Placopsis lateritioides I.M.Lamb

#### **FLORA CATEGORY**

Lichen - Native

#### **ENDEMIC TAXON**

Yes

#### **ENDEMIC GENUS**

Nο

### **ENDEMIC FAMILY**

Nο

#### STRUCTURAL CLASS

Lichens - Foliose

# **CURRENT CONSERVATION STATUS**

2018 | Not Threatened | Qualifiers: SO

#### **BRIEF DESCRIPTION**

Characterised by the rosette-forming to irregularly spreading, rustyorange-coloured thallus; and the grey-blue, scattered soralia.

#### **DISTRIBUTION**

**South Island**: Nelson (Mt Arthur, St Arnaud Range, Mt Lakeman), Canterbury (Phipps Peak, Temple Basin, Arthur's Pass, Craigieburn Range, Two Thumbs Range, Sebastopol, Mt Cook National Park, Mt Peel), Otago (Mt Ida, Swampy Hill, Flagstaff Hill Dunedin).

#### **HABITAT**

On exposed subalpine to high-alpine rocks, 650-2,000 m. Associating with *Placopsis bicolor*, *P. dennanensis*, *P. gelidioides*, *Umbilicaria cylindrica*.

# TEALAND YEALAND YEALAN



Photographer: Melissa Hutchison, Date taken: 09/01/2022, Licence: CC BY-NC.



Photographer: Melissa Hutchison, Date taken: 09/01/2022, Licence: CC BY-NC.

#### **DETAILED DESCRIPTION**

Thallus determinate, effigurate, orbicular to irregularly spreading, to 6 cm diam., saxicolous. Marginal lobes swollen, expanded at apices, rounded, crenate, discrete or becoming confluent, usually separated by narrow cracks, extending into centre of thallus, lobes also transversely cracked, wrinkled or folded. Upper surface distinctly rust-red to orange-brown, epruinose, areolate centrally, areolae swollen, rounded to irregular, 0.5-2 mm diam., isidia absent, sorediate. Soredia sparse to numerous, grey-white, erumpent, subrotund, to 2 mm diam., forming abraded patches on areolae. Cephalodia sparse, rounded to irregular, or radially lobate, 2-5 mm diam., brownish-red, dull, epruinose. Apothecia not seen. Pycnidia immersed in areolae, ostiole minute, punctiform. Conidia filiform, straight or curved 13-16 × 0.5 μm.

**Chemistry**: Thallus K-, C+ red, KC+ red, Pd-; containing gyrophoric acid.

#### **SUBSTRATE**

Saxicolous

# **ATTRIBUTION**

Fact sheet prepared by Melissa Hutchison (February 2023). Brief description, Distribution, Habitat, and Features sections copied from Galloway (1985, 2007).

# REFERENCES AND FURTHER READING

Galloway D.J. 1985: Flora of New Zealand: Lichens. Wellington: PD Hasselberg, Government Printer. 662 pp. Galloway D.J. 2007: Flora of New Zealand: Lichens, including lichen-forming and lichenicolous fungi. 2nd edition. Lincoln, Manaaki Whenua Press. 2261 pp.

# MORE INFORMATION

 $\underline{\text{https://www.nzpcn.org.nz/flora/species/placopsis-lateritioides/}}$