

LIST 1:SOME INDIGENOUS VASCULAR PLANTS IN BLOCK BETWEEN BANK ROAD AND ORANGI KAUPAPA ROAD, NORTHLAND, WELLINGTON, CENTRED ON NZMS 260 R27 MAP WELLINGTON, G.R. 577904; LIST COMPILED ON 29/8/96 BY B.MITCALFE AND J.C. HORNE. (NOTE : (A) PLANTED.

BOTANICAL NAME	MAAORI NAME	COMMON NAME
-----------------------	--------------------	--------------------

MONOCOT TREES

Cordyline australis

Rhopalostylis sapida (P)

DICOT TREES AND SHRUBS

Alectryon excelsus

Aristotelia serrata

Brachyglottis repanda

Coprosma grandifolia

Coprosma rhamnoides

Coprosma robust

Corokia cotoneaster (P)

Dodonea viscosa (P)

Elaeocarpus dentatus

Geniostoma rupestre

var. *ligustrifolium*

Hebe diosmifolia (P)

Hebe stricta var. *atkinsonii*

Hoheria populnea

Macropiper excelsum

Melicytus ramiflorus

Melicope ternata

Metrosideros excelsus

Myoporum laetum

Myrsine australis

Pittosporum ralphii

Pittosporum tenuifolium (P?)

Pseudopanax arboreus

Pseudopanax lessonii (garden escape)

Pseudopanax sp. (garden escape)

MONOCOT LIANES

Ripogonum scandens

DICOT LIANES

Muehlenbeckia australis

Parsonsia heterophylla

Passiflora tetrandra

Rubus cissoides

FERNS

Asplenium bulbiferum
Asplenium oblongifolium
Asplenium polyodon
Blechnum chambersii
Blechnum filiforme
Blechnum membranaceum
Cyathea dealbata
Cyathea medullaris
Histiopteris incisa
Hypolepis ambigua
Lastreopsis glabella
Lastreopsis velutina
Phymatosorus pustulatus
Polystichum richardii
Pteridium esculentum
Pyrrosia eleagnifolia

MONOCOT HERBS

Cortaderia sp.
Phormium cookianun (P?)

.....

LIST 2 - ADVENTIVE PLANTS.

Acer pseudoplatanus	sycamore
Agapanthus sp.	agapanthus
Banksia sp.	banksia
Buddleja davidii	buddleia
Cedronella caneriensis	balm of Gilead
Clematis vitalba	old man's beard
Cytisus scoparius	broom
Digitalis sp.	foxglove
Ehrharta erecta	veldt grass
Hedera helix	ivy
Pseudosuga menziesii	fir
Rubus fruticosus agg.	blackberry
Senecio mickanioides	German ivy
Senecio angulatus	Cape ivy
Solanum sp.	nightshade
Tradescantia fluminensis	wandering willie
Ulex europaeus	gorse

DESCRIPTION OF ORANGI KAUPAPA BLOCK.

ASPECT

SOUTHEAST

SLOPE

STEEP TO VERY STEEP

PHYSIOGRAPHY

GULLY AND SPUR

FOREST TYPE

MIXED INDIGENOUS 80%, AND EXOTIC - 20%, BROADLEAF.

80% CLOSED CANOPY

EXTENSIVE AREAS OF INDIGENOUS GROUND COVER (HOUND'S TONGUE FERN).

ONE LARGE, AND SEVERAL SAPLING WHARANGI TREES - (UNUSUAL).

MANY LARGE SYCAMORE.

SEVERAL LARGE *Pittosporum ralphii*, PROBABLY PLANTED.

ON BATTER IMMEDIATELY ABOVE NORTH SIDE OF BANK STREET PRIVATE DRIVE, VEGETATION HAS BEEN TRIMMED, INCLUDING A SUBSTANTIAL MAHOE, AND OCCASIONAL NATIVE TREES HAVE BEEN PLANTED.

IN GULLY BELOW MATAROA STREET, A CONSIDERABLE AMOUNT OF DOMESTIC RUBBISH HAD ACCUMULATED.