


(258)

HIGHER PLANTS (LYCOPDS, FERNS, GYMNOSPERMS, FLOWERING PLANTS) OF MOUNTAINS BETWEEN MATAKITAKI R. AND STATION CK, NORTH-WEST OF ALPINE FAULT, S. NELSON, 4000-4875 FT (1200-1500 M)

Highest point: unnamed peak, 4875 ft, S 39 828 378

A.P. Druce, Wellington, May 1984

(revised June 1986, August 1986, Nov. 1988, Sept. 1990, July 1991, Apr. 1992)  
(Based on visit March 1984)



(unc) - uncommon

- (F) - on non-calcareous sedimentary rocks (forest)
- (L) - on limestone ("Wooded Peak limestone") ( $\pm$  forest, burnt)
- (G) - on basic igneous rocks (forest/non-forest)
- (M) - on ultrabasic igneous rocks ("Mineral Belt") (non-forest)

INDIGENOUS SPECIES

GYMNOSPERM SHRUBS

- Lepidothamnus laxifolius* (G, M)
- Podocarpus nivalis* (G, unc M)

DICOT TREES

- Nothofagus menziesii* (F, L)
- N. solandri* var. *cliffortioides* (F, L, G)
- Pseudopanax colensoi* s.s. (unc F, unc G)

DICOT SHRUBS

- Brachyglottis bidwillii* var. *viridis* (unc G)
- Cassinia vauvilliersii* (unc G)

- Coprosma ciliata* ssp. *ciliata* (F)
- C. depressa* (F)
- C. perpusilla* ssp. *perpusilla* (unc M)
- C. pseudocuneata* ssp. (F, G)
- C. rugosa* ssp. *rugosa* (L)

- C. sp.* (unnamed; aff. *C. pseudo-cuneata*) (low-growing shrubs with small thick arcuate lvs) (G)
- Cyathodes dealbata* (incl. *C. pumila*) (G)
- Dracophyllum pronum* (M)
- D. uniflorum* (G, unc M)
- D. sp.* (unnamed sp; aff. *D. kirkii*) (low-growing shrubs; lf blade up to 2mm wide, slightly glaucous) (G, M)

*D. pronum* x *D. sp.* (unnamed) (M?)

*Exocarpus bidwillii* (G)

*Gaultheria crassa* (G)

*G. sp.* (unnamed; included in *G. depressa*, as var. *novae-zelandiae*, by Franklin 1962) (G)

*Hebe canterbutiensis* (G)

*H. ciliolata* (G)

*H. decumbens* (unc G)

*H. macrantha* var. *brachiphylla* (G)

*H. paucitamosa* (M)

*Helichrysum intermedium* var. (G)

*Melicytus alpinus* s.s. (G, M)

*Leucopogon fraseri* s.s. (L, G, M)

*L. suaveolens* (G, M)

*Myrsine divaricata* s.s. (F, G)

*M. nummularia* (G, unc M)

*Olearia lacunosa* var. *lacunosa* (F)

↳ (include *Cyathodes colensoi*)

## GRASSES

*O. nummulariifolia* var. (lvs  
distinctly longer than broad,  
rather than suborbicular as  
in var. *nummulariifolia*) (unc G)

*Pentachondra pumila* (G, M)

*Pimelea suteri* var. (*P. oreophila*)  
(G, M)

*Agrostis personata* (L)

*A. sp.* (unnamed; aff. *A.*

*dyeri*) (panicle spreading, not  
contracted as in *A. dyeri*) (G,  
unc M)

*Chionochloa australis* (G)

*C. conspicua* ssp. *conspicua* (F, L)

*C. defracta* (M)

*C. pallens* ssp. *pilosa* (L)

*C. sp.* (unnamed) (*C. flav-*  
*escens* auct. S. L., non *C. flav-*  
*escens* Zotor) (unc G)

## DICOT LIANES AND RELATED

### TRAILING PLANTS

*Muehlenbeckia axillaris* (G)

## LYCOPPODS

*Lycopodium australianum* (G, unc M)

*L. fastigiatum* (F, L, G)

## FERNS

*Blechnum penna-matina* (L, unc M)

*Cystopteris tasmanica* (L)

*Grammitis poeppigiana* (unc M) *Deyeuxia avenoides* (incl. *D. a.*

*Hypolepis millefolium* (F, L)

*var. brachyantha* (L)

*Ophioglossum coriaceum* agg. (L)

*Festuca* sp. (*F. matthewsii*

*Polystichum vestitum* (F, L)

*sensu* Connor 1968) (M)

*Hierochloe* sp. (unc M)

*Koeleria* sp. (*K. novozelandica* agg.)  
(lvs up to 3mm wide, flat) (G, unc M)

*Lachnagrostis* sp. (unnamed;  
included in *L. filiformis*, as  
var. *semiglabra*, by Zotor  
1965) (L, unc M)

## ORCHIDS

*Chiloglottis cornuta* (F, L)

*Corybas trilobus* s.s. (F)

*Piasophyllum colensoi* (G)

*Poa breviglumis* (F, L, unc M)

*P. colensoi* (L, G, M)

*P. imbecilla* (L)

*P. kirkii* (L)

*P. maia* (F)

*P. sp.* (unnamed; aff. *P. novae-zelandiae*) (G)

*Rytidosperma setifolium* (G, M)

*Trisetum* sp. (unnamed) (plants slender; lf sheath glabrous or nearly so; lf blade flat, up to 2.5 cm wide, margins smooth, vs. with a few hairs near ligule) (F, L)

#### SEDGES

*Carex aciculatis* (G)

*C. breviculmis* (L, G)

*C. comans* (L)

*C. wakatipu* s.s. (L)

*Carpha alpina* s.s. (M)

*Oreobolus pectinatus* (M)

*Schoenus pauciflorus* s.s. (G, M)

*Uncinia caespitosa* (G)

*U. clavata* (F)

*U. divaricata* (L, G, M)

*U. filiformis* (F)

*U. fuscovaginata* var. (*glaucus*) (L)

*U. nervosa* s.s. (G)

*U. rupestris* (incl. *U. angustifolia*) (F, G)

*U. sp.* (unnamed) (*U. affinis* sensu Hamlin 1959, non *U. riparia* var. *affinis* Col. ex Clarke) (L)

#### RUSHES

*Luzula* sp. (i) (unnamed; aff. *L. tufa*)\* (G, M)

*L. sp.* (ii) (unnamed; included in *L. picta*, as var. *limosa*, by Edgar 1966) (F)

#### MONOCOT HERBS OTHER THAN ORCHIDS, GRASSES, SEDGES, RUSHES

*Astelia nervosa* s.s. (G)

*Phormium cookianum* (G)

#### COMPOSITE HERBS

*Anaphalis* sp. (*Helichrysum bellidoides* s.s.) (L, G, M)

*Brachyglottis* sp. (unnamed; included in *B. bellidoides*, as var. *crassa*, by Nordenstam 1978) (G)

*Brachyscome radicata* (L)

*B. sinclairii* s.s. (G)

*Celmisia gracilenta* ssp. (G)

*C. incana* (incl. *C. discolor*) (G)

*C. latifolia* (G)

*C. montoi* ssp. (*C. semicordata* ssp. *semicordata*) (G, unc M)

*C. sinclairii* (incl. *C. allanii*, *C. dumicola*) (G)

*C. spectabilis* ssp. (G)

\* (lvs up to 2.5 cm wide, channelled, strict; heads 6-9-(12) mm diam. tepals slightly > capsules)

*Craspedia* sp. (i) (unnamed) (lvs green, both surfaces clad in multicellular non-glandular hairs; margins faintly to distinctly white with cottony hairs; fls white) (unc M)

*C. sp. (ii)* (unnamed; included in *C. lanata*, as var. *elongata*, by Allan 1961) (lvs white or greenish white, both surfaces ± clad in cottony hairs; fls white) (G, M)

*C. sp. (iii)* (lvs green, almost glabrous) (unc M)

*Gnaphalium tuahinicum* (L)

*Haastia sinclairii* var. *sinclairii* (unc G)

*H. filicawle* (L)

*Lagenifera pinnatifida* (F)

*L. pumila* (L)

*L. strangulata* (F, L, G)

*Leptinella* sp. +

*Leucogenes grandiceps* (G)

*Raoulia apicinigra* s.s. (G)

*R. bryoides* (G)

*R. glabra* (L)

*R. grandiflora* var. *grandiflora* (G)

*R. tenuecaulis* (incl. *R. t.* var. *dimorpha* and *R. t.* var. *pusilla*) (L)

*Taraxacum magellanicum* (L)

**DICOT HERBS OTHER THAN  
COMPOSITES**

*Acaena profundeincisa* (L)

*Aciphylla montoi* (G)

*A. sp. (unnamed; aff. A. aurea)* (stipules short, spreading) (G)

*Anisotome aromatica* (G, unc M)

*A. filifolia* (unc M)

*A. haastii* (G)

*A. pilifera* (unc G)

*Colobanthus affinis* (L)

*C. sp. (unnamed)* (resembling *Sagina procumbens*) (unc G, M)

*Epilobium alsinoides* s.s. (L)

*E. atriplicifolium* s.s. (G)

*E. brunnescens* s.s. (L)

*E. glabellum* s.s. (G)

*E. Komarovianum* (L)

*E. tenuipes* s.s. (L)

*E. pycnostachyum* (G)

*E. sp. (unnamed; included in*

*E. brunnescens*, as ssp. *minutiflorum*, by Raven and Engelhorn, 1971) (L, unc M)

*Euphrasia montoi* (G)

*Forstera tenella* (G)

*Gingidia decipiens* (G)

*Gentiana* sp. (cf. *G. corymbifera*)

*G. montana* (incl. *G. montana* var. *stolonifera*, *G. patula*, and *G. bellidifolia* var. *pulchella*)

+ (included in *H. squamula*, as *ssp. mediaea*, by Lloyd and Webb, 1987)

*Hydrocotyle microphylla* (L)  
H.sp. (unnamed; included in *H. novae-zealandiae*, as var. *montana*, by Kirk 1899) (L)

*Myosotis traversii* s.s. (unc G)  
*Notothlaspi* sp. (unnamed) (*N. australis* auct., non *Thlaspi australe* Hook.f.)  
*Oreomytthis colensoi* s.s. (incl. *O.c.* var. *hispida* and *O.c.* multi-  
fida) (M)

*O. sp.* (unnamed; aff. *O. ramosa*) (FL)  
*Ourisia lactea* ssp. *lactea* (L)

*Oxalis magellanica* (FL)  
*Patahebe lyallii* (unc M)  
*Ranunculus membranifolius* (FL)  
*R. reflexus* (unc L)

*Schizilema roughii* (G)  
*Scleranthus brockiei* (unc M)  
*Stackhousia minima* (L, G)  
*Urtica incisa* (L)  
*Viola cunninghamii* (unc M)  
*Wahlenbergia albomarginata*  
var. *albomarginata* (incl. *W. pygmaea* var. *laxa*, *W. brockiei*,  
and *W. simpsonii*) (L, G, M)

## ADVENTIVE SPECIES

### GRASSES

*Agrostis capillaris* (L)

### COMPOSITE HERBS

*Cirsium atropurpureum* (L)

*C. vulgare* (L)

*Hieracium pilosella* (L)

*Hypochoeris radicata* (L, G)

*Mycelis muralis* (L, G)

*Taraxacum officinale* s.l. (L)

### DICOT HERBS OTHER THAN COMPOSITES

*Cerastium fontanum* ssp.  
*vulgare* (L)

*Origanum vulgare* (unc, L)

*Rumex acetosella* (L)

*Sagina procumbens* (L)

*Trifolium repens* (L)

No. of adventive species: 12.

No. of spp. in different areas:

(F) (non-calc. sed. rocks) : 26 I ; nil A

(L) (limestone) : 54 I ; 12 A

(G) (basic igneous rocks) : 84 I ; 2 A

(M) (ultrabasic " " ) : 47 I ; nil A

I - indigenous spp

A - adventive spp.

No. of indigenous species: 152.

## REFERENCES

- Allan, H.H. 1961: "Flora of New Zealand". Government Printer, Wellington. Vol. I. 1085 pp.
- Connor, H.E. 1968: Interspecific hybrids in hexaploid New Zealand Festuca. N.Z. Jl Bot. 6: 295-308.
- Edgar, E. 1966: Luzula in New Zealand. N.Z. Jl Bot. 4: 159-84.
- Franklin, D.A. 1962: The Ericaceae in New Zealand (Gaultheria and Pernettya). Trans. R. Soc. N.Z. Bot. 1: 155-73.
- Hamlin, B.G. 1959: A revision of the genus Uncinia (Cyperaceae - Carexoidae) in New Zealand. Dom. Mus. Bull. 19. 106 pp.
- Kirk, T. 1899: "The Student's Flora of New Zealand." Government Printer, Wellington.
- Lloyd, D.G.; Webb, C.J. 1987. The reinstatement of Leptinella at generic rank, & the status of the 'Cotulaceae' (Asteraceae, Anthemideae). N.Z. Jl Bot. 25: 99-105.
- Raven, P.H.; Engelhorn, T. 1971: New taxa and new combinations in Australasian Epilobium (Onagraceae). N.Z. Jl Bot. 9: 345-50.
- Zotov, V.D. 1965: Grasses of the subantarctic islands of the New Zealand region. Rec. Dom. Mus. Wellington 5: 101-46.
- Nordanstam, B. 1978: Opera Bot. 44: 1-84.
-

