

g. 2000

List of Plants Horohoro Stream to Belmont Trig.

7 May 1949.

Tall forest at lower levels. Scrub on summit

Ferns

- | | |
|--------------------------|---|
| Polypodium diversifolium | Asplenium floccidum |
| Blechnum lauceolatum | Leptopteris hymenophylloides |
| B. fluviatile | Asplenium bulbiferum |
| Dryopteris penigera | A. lucidum |
| Polypodium pustulatum | A. adiantoides |
| Polystichum richardi | Histiopteris incisa |
| P. hispidum | Blechnum filifolium ^{crane} crane |
| Trichomanes venosum | Cyathea medullaris |
| Blechnum discolor | Cy. dealbata |
| Hemitelia smithii | Dicksonia squarrosa |
| Polystichum adiantiforme | Cardiomanes reniforme |
| Pteris maileuta | Pacria scaberula |
| Polystichum vestitum | |

Fern allies

- Lycopodium billardieri

Tall Trees

- | | |
|--------------------------|--------------------------|
| Beilschmiedia tawa | Knightia exeelsa |
| Laurelia novaezealandiae | Cuscutina littoralis |
| Hedycarya arborea | Podocarpus totara |
| Podocarpus ferruginea | Elaeocarpus dentatus |
| P. dactyloides | X Corynocarpus laevigata |
| Dacrydium cupressinum | |
| X Dypsoxylum spectabile | |

Small Trees

- | | |
|----------------------------------|--------------------------------|
| X <i>Hebe parviflora</i> | <i>Hebe salicifolia</i> |
| X <i>Oleaia paniculata</i> | <i>Pittosporum tenuifolium</i> |
| <i>Brachyglottis repanda</i> | <i>Alectryon excelsum</i> |
| <i>Euttonia australis</i> | <i>Pennantia conyubora</i> |
| <i>Coprosma robusta</i> | <i>Nothopanax arboreum</i> |
| <i>Schefflera digitata</i> | <i>Fuchsia exorticata</i> |
| <i>Aristotilia serrata</i> | X <i>Melicope ternata</i> |
| <i>Geniostoma ligustifolium</i> | <i>Myrtus bullata</i> |
| <i>Coprosma australis</i> | <i>Oleaia rani</i> |
| X <i>C. areolata</i> | <i>Carpodetus serratus</i> |
| <i>Pittosporum eugenioides</i> | <i>Euttonia salicina</i> |
| <i>Pseudopanax crassifolium</i> | <i>Olea lanceolata</i> |
| { <i>Pseudowintera axillaris</i> | X <i>Coprosma foetidissima</i> |
| { <i>P. colorata</i> | <i>Nothopanax anomalum</i> |
| <i>Rhopalostylis sapida</i> | |

Shrubs

- | | |
|-----------------------------|---|
| <i>Coprosma rhamnoides</i> | |
| <i>Cassinia leptophylla</i> | } probably introduced since forest was cleared. |
| <i>Oleaia solandri</i> | |
| <i>Macropiper excelsum</i> | |
| <i>Senecio Kirkii</i> | |
| <i>Solanum aviculare</i> | |
| <i>Urtica ferox</i> | |

Lianes

Metroideros scandens

M. perforatum

Tetrapathaea tetrandra

Rubus cirroides (^{australis} _{aut.})

Rhipogonum scandens

Freycineta banksii

Parsonia heterophylla

Metroideros diffusa

Epiphytes

Artelia splendens

Collosporum hastatum

Griechina lucida

Earina mucronata

E. autumnalis

Herbaceous plants.

Libertia isoioides

Uncinia uncinata

Microlaena avenacea

Helichrysum bellidioides

Erechtites prenanthoides

The following formed the wind swept scrub on the summit of Belmont Trig.

Pseudowintera colorata

Nothofagus anomalum

Coprosma foetidissima

Coprosma rhamnoides

Carpodetus serratus

Metroideros diffusa

Polyptichum vestitum

Paesia scaberula

Blechnum discolor

